

An Introduction to Modern Astrophysics notes

PB21151823 杨景硕

Peculiar Motions and the Local Standard of Rest

The velocity components of stars in the solar neighborhood are traditionally labeled

$$\Pi \equiv \frac{dR}{dt}, \Theta \equiv R \frac{d\theta}{dt}, Z \equiv \frac{dz}{dt}$$

Defining the **dynamical local standard of rest** (dynamical LSR) to be a point that is instantaneously centered on the Sun and moving in a perfectly circular orbit along the solar circle about the Galactic center.

$$\Pi_{LSR} = 0, \Theta_{LSR} = \Theta_0, Z_{LSR} = 0$$

An alternative definition for the LSR known as the **kinematic local standard of rest** (kinematic LSR) is based on the average motions of stars in the solar neighborhood.

$$\langle \Pi \rangle = 0, \langle \Theta \rangle = \Theta_0, \langle Z \rangle = 0$$

The velocity of a star relative to the dynamical LSR is known as the star's **peculiar velocity** and is given by

$$u = \Pi - \Pi_{LSR} = \Pi, v = \Theta - \Theta_{LSR} = \Theta - \Theta_0, w = Z - Z_{LSR} = Z$$

The average of u , v and w for all stars in the solar neighborhood, excluding the Sun, is

$$\langle u \rangle = 0, \langle v \rangle = C\sigma_u^2 = C \langle u^2 \rangle < 0, \langle w \rangle = 0$$

Why $\langle \Theta \rangle < \Theta_0$ ($\langle v \rangle < 0$)?

- The stars inside the Sun's orbit are in the **apogalacticon** of their elliptical orbits when they pass through the Sun's neighborhood, so the tangential velocity Θ is smaller.
- The stars outside the Sun's orbit are in the **perigalacticon** of their elliptical orbits when they pass through the Sun's neighborhood, so the tangential velocity Θ is larger.
- There are more stars inside the Sun's orbit than beyond it.

Why should σ_u correlate with $\langle v \rangle$?

- larger σ_u , wider range of elliptical orbits included, more negative $\langle v \rangle$
- smaller σ_u , fewer stars with orbits noncircular, $\langle v \rangle \sim 0$

Differential Galactic Rotation and Oort's Constants

The relative radial and transverse velocities of a star (at point S) to the Sun (at point O) are, respectively,

$$v_r = \Theta \cos \alpha - \Theta_0 \sin l$$
$$v_t = \Theta \sin \alpha - \Theta_0 \cos l$$




FIGURE 22 The geometry of analyzing differential rotation in the Galactic plane. The Sun is at point O , the center of the Galaxy is located at C , and the star is at S , located a distance d from the Sun. ℓ is the Galactic longitude of the star at S , and α and β are auxiliary angles. The directions of motion reflect the clockwise rotation of the Galaxy as viewed from the NGP.

where Θ_0 is the orbital velocity of the Sun in the idealized case of perfectly circular motion (actually the orbital velocity of the LSR) and α is defined in the figure. Defining the *angular-velocity curve* to be

$$\Omega(R) \equiv \frac{\Theta(R)}{R},$$

the relative radial and transverse velocities become

$$v_r = \Omega R \cos \alpha - \Omega_0 R_0 \sin \ell,$$

$$v_t = \Omega R \sin \alpha - \Omega_0 R_0 \cos \ell.$$

Now, by referring to the geometry of Fig. 22 and considering the right triangle ΔOTC , we find

$$R \cos \alpha = R_0 \sin \ell,$$

$$R \sin \alpha = R_0 \cos \ell - d.$$

Substituting these relations into the previous expressions, we have

$$v_r = (\Omega - \Omega_0) R_0 \sin \ell, \quad (37)$$

$$v_t = (\Omega - \Omega_0) R_0 \cos \ell - \Omega d. \quad (38)$$

Equations (37) and (38) are valid as long as the assumption of circular motion is justified.

Oort derived a set of approximate equations for v_r and v_t that are valid only in the region near the Sun.

Defining the **Oort constants**

$$A \equiv -\frac{1}{2} \left(\frac{d\Theta}{dR} \Big|_{R_0} - \frac{\Theta_0}{R_0} \right)$$

$$B \equiv -\frac{1}{2} \left(\frac{d\Theta}{dR} \Big|_{R_0} + \frac{\Theta_0}{R_0} \right)$$

Using the Taylor expansion of $\Omega(R)$, the difference between Ω and Ω_0 , and the approximate value of Ω is

$$\Omega - \Omega_0 \approx \frac{d\Omega}{dR} \Big|_{R_0} (R - R_0) = \left(\frac{1}{R_0} \frac{d\Theta}{dR} \Big|_{R_0} - \frac{\Theta_0}{R_0^2} \right) (R - R_0) = (-2A/R_0)(-d\cos l)$$

$$\Omega \approx \Omega_0 = A - B$$

Inserting equations above into Eqs. (37) and (38) results in

$$v_r \approx Ad \sin 2l$$

$$v_t \approx Ad \cos 2l + Bd$$

Nine Equations of Cosmology

$$\text{Friedmann equation : } H^2(t) \equiv \left(\frac{1}{R(t)} \frac{dR(t)}{dt} \right)^2 = \frac{8\pi G}{3} \rho(t) - \frac{kc^2}{R^2(t)} = \frac{8\pi G}{3} \rho_c(t)$$

$$\text{where } \rho(t) = \rho_m(t) + \rho_r(t) + \rho_\Lambda, \rho_\Lambda = \frac{\Lambda c^2}{8\pi G} = \text{const}$$

$$\text{Density parameter : } \Omega(t) \equiv \frac{\rho(t)}{\rho_c(t)} = 1 - \frac{kc^2}{R^2(t)H^2(t)}$$

$$\text{where } \Omega(t) = \Omega_m(t) + \Omega_r(t) + \Omega_\Lambda(t), \Omega_\Lambda(t) = \frac{\Lambda c^2}{3H^2(t)}$$

$$\text{Fluid equation : } \frac{d(R^3(t)\rho(t))}{dt} = -\frac{P(t)}{c^2} \frac{dR^3(t)}{dt}$$

$$\text{where } P(t) = P_m(t) + P_r(t) + P_\Lambda$$

$$\text{Acceleration equation : } \frac{d^2 R(t)}{dt^2} = -\frac{4\pi G}{3} \left(\rho(t) + \frac{3P(t)}{c^2} \right) R(t)$$

$$\text{Equation of state : } P(t) = w\rho(t)c^2$$

$$\text{where } w_m = 0, w_r = \frac{1}{3}, w_\Lambda = -1$$

$$\text{Cosmological redshift : } 1 + z = \frac{\lambda_0}{\lambda_e} = \frac{R(t_0)}{R(t)} = \frac{1}{R(t)}$$

$$\text{Density evolution equation : } R^{3(1+w)}(t)\rho(t) = \rho_0 \text{ or } \rho(z) = \rho_0(1+z)^{3(1+w)}$$

$$\text{Hubble parameter evolution equation : } H^2(z) = H_0^2(\Omega_{m,0}(1+z)^3 + \Omega_{r,0}(1+z)^4 + \Omega_{\Lambda,0} + (1-\Omega_0)(1+z)^2)$$

where the subscript 0 represents the present value

$$\text{Deceleration parameter : } q(t) \equiv -\frac{R(t)}{(dR(t)/dt)^2} \frac{d^2 R(t)}{dt^2} = -\frac{1}{R(t)H^2(t)} \frac{d^2 R(t)}{dt^2} = \frac{1}{2}(1+3w_i)\Omega_i(t)$$

The constant k determines the ultimate fate of the universe:

- If $k > 0$, the total energy of the shell is negative, and the universe is *bounded*, or **closed**. In this case, the expansion will someday halt and reverse itself.
- If $k < 0$, the total energy of the shell is positive, and the universe is *unbounded*, or **open**. In this case, the expansion will continue forever.
- If $k = 0$, the total energy of the shell is zero, and the universe is **flat**, neither open nor closed. In this case, the expansion will continue to slow down, coming to a halt only as $t \rightarrow \infty$ and the universe is infinitely dispersed.

Transition from the Radiation Era to the Matter Era to the Λ Era

The behavior of the scale factor $R(t)$ for a flat universe can be found by setting $k = 0$ in the Friedmann equation.

Radiation era when $t \ll t_{r,m}$: $R(t) \propto t^{1/2}$

The transition from the radiation era to the matter era occurred when

$$R_{r,m} \approx 3.05 \times 10^{-4}, z_{r,m} \approx 3270, t_{r,m} \approx 5.52 \times 10^4 \text{ yr}$$

Matter era when $t_{r,m} \ll t \ll t_H$: $R(t) \propto t^{2/3}$

The acceleration of the universe changed sign (from negative to positive) when

$$R_{accel} \approx 0.57, z_{accel} \approx 0.76, t_{accel} \approx 7.08 \text{ Gyr}$$

The transition from the matter era to the Λ era occurred when $R_{m,\Lambda} \approx 0.72, z_{m,\Lambda} \approx 0.39, t_{m,\Lambda} \approx 9.55 \text{ Gyr}$

Λ era when $t \gg t_H$: $R(t) \propto e^{H_0 t \sqrt{\Omega_{\Lambda,0}}}$

Distances to the Most Remote Objects in the Universe

The **Robertson-Walker metric** determines the spacetime interval between two events in an isotropic, homogeneous universes.

$$ds^2 = c^2 dt^2 - R^2(t) \left(\frac{d\varpi^2}{1 - k\varpi^2} + \varpi^2 d\theta^2 + \varpi^2 \sin^2 \theta d\phi^2 \right)$$

With $ds = 0$ for a light ray, and $d\theta = d\phi = 0$ for a radial path traveled from the point of the light's emission at **comoving coordinate** ϖ_e to its arrival at Earth at $\varpi = 0$, taking the negative square root (so ϖ decreases with increasing time) gives

$$\frac{-cdt}{R(t)} = \frac{d\varpi}{\sqrt{1 - k\varpi^2}}$$

$$\therefore \int_{t_e}^{t_0} \frac{cdt}{R(t)} = \int_0^{\varpi_e} \frac{d\varpi}{\sqrt{1 - k\varpi^2}} = \begin{cases} \varpi_e \dots \text{for } \Omega_0 = 1, \text{ then } k = 0 \\ \sin^{-1} \varpi_e \dots \text{for } \Omega_0 > 1, \text{ then } k > 0 \\ \sinh^{-1} \varpi_e \dots \text{for } \Omega_0 < 1, \text{ then } k < 0 \end{cases}$$

Defining two dimensionless integrals

$$I(z) \equiv H_0 \int_{t_e}^{t_0} \frac{dt}{R(t)} = H_0 \int_0^z \frac{dz'}{H(z')} = z - \frac{1}{2}(1 + q_0)z^2 + \left(\frac{1}{6} + \frac{2}{3}q_0 + \frac{1}{2}q_0^2 + \frac{1}{6}(1 - \Omega_0)\right)z^3 - \dots$$

where $\frac{dz}{dt} = -\frac{1}{R^2(t)} \frac{dR(t)}{dt} = -\frac{H(t)}{R(t)}$

$$S(z) \equiv \begin{cases} I(z) \dots \text{for } \Omega_0 = 1, \text{ then } k = 0 \\ \frac{1}{\sqrt{\Omega_0 - 1}} \sin(I(z)\sqrt{\Omega_0 - 1}) \dots \text{for } \Omega_0 > 1, \text{ then } k > 0 \\ \frac{1}{\sqrt{1 - \Omega_0}} \sinh(I(z)\sqrt{1 - \Omega_0}) \dots \text{for } \Omega_0 < 1, \text{ then } k < 0 \end{cases} \approx z - \frac{1}{2}(1 + q_0)z^2 \dots \text{for } z \ll 1$$

Therefore the comoving coordinate as a function of the redshift is $\varpi(z) = \frac{c}{H_0} S(z)$. Now we are ready for the concept of four distances at time t_0 .

$$\text{Coordinate distance : } r_0(z) = \varpi(z) = \frac{c}{H_0} S(z)$$

$$\text{Proper distance : } d_{p,0}(z) \equiv \int_{t_e}^{t_0} \frac{cdt}{R(t)} = \frac{c}{H_0} I(z)$$

$$\text{Luminosity distance : } d_{L,0}(z) \equiv r_0(z)(1 + z) = \frac{c}{H_0} S(z)(1 + z)$$

$$\text{Angular diameter distance : } d_{A,0}(z) \equiv \frac{r_0(z)}{1 + z} = \frac{c}{H_0} \frac{S(z)}{1 + z}$$

Multiplying by the scale factor $R(t)$ then converts these to the distances at some other time t .